



**Public Reporting for 2017Q1 - 2017Q4  
CathPCI Registry®**

**Hospital Performance: 731406 - Cooper University Health System compared to all eligible registry hospitals**

**DQR Status as of Aggregation Date:**

Q1	Q2	Q3	Q4

**CathPCI Public Reporting:** Participating with ACC

**Publication Status:** Report available for Publication

**Aggregation Date:** Apr 17, 2018 11:59:59 PM

Metric ID	Cardiosmart Metric Description	Rating	Detailed Score	95% Interval Estimate
0	<b>Number of PCI/angioplasty procedures performed during the calendar year.</b> The number of PCI/angioplasty procedures a site performs does not necessarily indicate higher quality, but it may be an indication of how experienced this site is with the procedure.	1,016 Procedures	Reporting for 4 Quarters	
8	<b>Use of Aspirin to reduce the chance of blood clots after PCI/angioplasty.</b> Patients should be prescribed Aspirin to reduce the risk of heart attacks caused by blood clots in new stents after having a PCI/angioplasty- unless there is a reason not to use the medicine (such as an allergy).  This score shows how well this facility is following this guideline - higher is better. Patients who cannot take aspirin are excluded.	★★★★★	P = 99.70% 95% IE = (99.58%,99.82%)	
		* As data displayed for public reporting participating hospitals on <a href="https://www.cardiosmart.org/Resources/Find-Your-Heart-a-Home">https://www.cardiosmart.org/Resources/Find-Your-Heart-a-Home</a>		



**Public Reporting for 2017Q1 - 2017Q4  
CathPCI Registry®**

**Hospital Performance: 731406 - Cooper University Health System compared to all eligible registry hospitals**

DQR Status as of Aggregation Date:

Q1	Q2	Q3	Q4

**CathPCI Public Reporting:** Participating with ACC

**Publication Status:** Report available for Publication

**Aggregation Date:** Apr 17, 2018 11:59:59 PM

Metric ID	Cardiosmart Metric Description	Rating	Detailed Score	95% Interval Estimate
9	<p><b>Use of a P2Y12 inhibitor medication to reduce the chance of blood clots after PCI/angioplasty.</b> Patients should be prescribed a P2Y12 inhibitor medication to reduce the risk of heart attacks caused by blood clots in new stents after having a PCI/angioplasty- unless there is a reason not to use the medicine (such as an allergy).</p> <p>This score shows how well this facility is following this guideline - higher is better. Patients who cannot take P2Y12 inhibitor medicines are excluded.</p>	★★★★★	P = 99.87% 95% IE = (99.81%,99.92%)	
10	<p><b>Use of a Statin to decrease cholesterol after PCI/angioplasty.</b> Patients should be prescribed a Statin to decrease cholesterol and reduce the risk of heart attacks after having a PCI/angioplasty- unless there is a reason not to use the medicine (such as an allergy).</p> <p>This score shows how well this facility is following this guideline - higher is better. Patients who cannot take Statin medications are excluded.</p>	★★★★★	P = 99.25% 95% IE = (98.94%,99.56%)	
<p>* As data displayed for public reporting participating hospitals on <a href="https://www.cardiosmart.org/Resources/Find-Your-Heart-a-Home">https://www.cardiosmart.org/Resources/Find-Your-Heart-a-Home</a></p>				



**Public Reporting for 2017Q1 - 2017Q4  
CathPCI Registry®**

**Hospital Performance: 731406 - Cooper University Health System compared to all eligible registry hospitals**

DQR Status as of Aggregation Date:

Q1	Q2	Q3	Q4

**CathPCI Public Reporting:** Participating with ACC

**Publication Status:** Report available for Publication

**Aggregation Date:** Apr 17, 2018 11:59:59 PM

Metric ID	Cardiosmart Metric Description	Rating	Detailed Score	95% Interval Estimate
38	<p><b>Use of all recommended medications (Aspirin, P2Y12 inhibitor medication, and Statin) to reduce the chance of blood clots and decrease cholesterol after PCI/angioplasty.</b> Patients should be prescribed Aspirin, a P2Y12 inhibitor medication, and a Statin medication after having a PCI/angioplasty to reduce the chance of blood clots in new stents ,decrease cholesterol and reduce the risk of heart attacks- unless there is a reason not to use these medicines (such as an allergy).</p> <p>This score shows how well this facility is following this guideline - higher is better. Patients who cannot take all of the recommended medicines are excluded.</p>	★★★★★	P = 98.92% 95% IE = (98.46%,99.36%)	
		<p>* As data displayed for public reporting participating hospitals on <a href="https://www.cardiosmart.org/Resources/Find-Your-Heart-a-Home">https://www.cardiosmart.org/Resources/Find-Your-Heart-a-Home</a></p>		